



HUSKY® 800 NEUTRAL DISINFECTANT CLEANER

SPECIFICATIONS:

| | |
|---------------|-----------------------------|
| Registrations | EPA Registered Disinfectant |
| EPA# | 47371-131-8155 |
| Dilution | 1:64 |
| pH Level | 7.2 – 8.2 |
| Color | Blue |
| Fragrance | Ocean Breeze |

This **ONE-STEP** cleaner, disinfectant and deodorizer is ideal for use in hospitals, nursing homes, veterinary clinics, schools and federally inspected meat and poultry establishments.



Neutral, quaternary-based disinfectant cleaner for use on a wide variety of hard non-porous surfaces. This product is ideal for use on finished floors due to its neutral pH. It inhibits the growth of mold and mildew and controls related odors. EPA registered as a broad spectrum disinfectant cleaning product.

FORMULA FEATURES:

- Cleans, Disinfects and Deodorizes in One-Step
- Neutral pH
- Inhibits Mold Growth
- Effective against SARS-COV-2 in 1 minute⁸
- Hospital Use Disinfecting
- Powerful Fragrance Leaves Area Smelling Fresh and Clean
- Bloodborne Pathogen Compliant

⁸hard, non-porous surfaces

RECOMMENDED FOR:

- Finished Floors
- Walls
- Metal Surfaces
- Stainless Steel
- Glazed Porcelain
- Plastic Surfaces
- Toilet Bowl Surfaces and Urinals
- Bathroom Fixtures
- Bathtubs and Shower Stalls

DIRECTIONS:

DISINFECTION DILUTION: 1:64 (660 ppm quat) 2 ounces per gallon of water

This product is not for use on medical device surfaces.

DISINFECTION / CLEANING / DEODORIZING DIRECTIONS: Remove heavy soil deposits from surface, then thoroughly wet surface with a use-solution of 2 ounces of the concentrate per gallon of water. The use-solution can be applied with a cloth, mop, sponge, or coarse spray or by soaking. For sprayer applications, use a coarse spray device. Spray 6-8 inches from the surface, rub with a brush, cloth or sponge. Do not breathe spray. Let solution remain on surface for a minimum of 10 minutes. Rinse or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Food contact surfaces must be thoroughly rinsed with potable water. This product must not be used to clean the following food contact surfaces: utensils, glassware and dishes.

NEUTRAL DISINFECTANT CLEANER

DIRECTIONS: (CONTINUED)

TOILET BOWLS: Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying use-solution around the bowl and up under the rim. Stubborn stains may require brushing. To disinfect, first pre-clean to remove heavy soil, then remove or expel over the inner trap the residual bowl water. Pour in three ounces of the diluted solution. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush.

FUNGICIDAL DIRECTIONS: For use in areas such as locker rooms, dressing rooms, shower and bath areas, and exercise facilities follow disinfection directions.

MILDEWSTATIC INSTRUCTIONS: Will effectively control the growth of mold and mildew, plus the odors caused by them when applied to hard, nonporous surfaces such as walls, floors, and table tops. Apply use-solution of 2 ounces per gallon of water with a cloth, mop, sponge or coarse spray. Make sure to wet all surfaces completely. Let air dry. Repeat application weekly or when growth reappears.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS: For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2 oz. of this product per gallon of water for a period of 10 minutes. (Use 8 oz per gallon of water to kill Adenovirus Type 7.) Immerse all animal handling and restraining equipment as well as forks, shovels and scrapers used to remove litter and manure. Wipe or allow to air dry. Ventilate buildings and other closed spaces. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried. For Heavy Duty Cleaning and/or Odor Control: When greater cleaning is desired, use 4-8 oz of this product per gallon of water. Heavily soiled areas may require repeated cleaning before treatment.

DISINFECTANT CLAIMS:

- Pseudomonas aeruginosa
- Staphylococcus aureus
- Salmonella enterica
- Enterococcus faecalis - Vancomycin Resistant (VRE)
- Listeria monocytogenes
- Streptococcus pyogenes (Strep)
- Staphylococcus aureus - Community Associated Methicillin - Resistant (CA-MRSA)
- Staphylococcus aureus - Methicillin Resistant (MRSA)
- Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA)
- Enterobacter aerogenes
- Enterobacter cloacae - NDM-1
- Escherichia coli

VIRUCIDAL ACTIVITY:

- Hepatitis B Virus (HBV)
- Hepatitis C Virus (HCV)
- Herpes Simplex Virus Type 1
- Herpes Simplex Virus Type 2
- HIV-1 (AIDS virus)
- Influenza A Virus / Hong Kong
- Rotavirus (WA)
- Rubella (German Measles)
- SARS Associated Coronavirus
- SARS -Related Coronavirus 2 (SARS-CoV-2)

FUNGICIDAL ACTIVITY:

- Aspergillus niger
- Candida albicans
- Trichophyton mentagrophytes (Athlete's Foot Fungus)

ANIMAL VIRUSES:

- Avian Influenza H5N1
- Avian Influenza H5N1
- Canine distemper
- Equine Herpes Virus Type 1
- Feline coronavirus
- Feline leukemia
- Feline panleukopenia virus
- Infectious bovine rhinotracheitis
- Porcine parvovirus
- Pseudorabies (PRV)
- Transmissible Gastroenteritis virus (TGE)

This is not a complete list of organisms and names